

Prosma[®]

Presentation

Prosma[®] Eye Drop: A clear colourless, isotonic solution that is sterile until the bottle is opened. Each ml contains Ketotifen 0.25mg as Fumarate BP.

Description

Prosma[®] is a non-bronchodilator anti-asthmatic drug with marked anti-anaphylactic property. In addition, Prosma[®] exerts a powerful and sustained H₁ receptor blocking activity which can be clearly dissociated from its anti-anaphylactic property.

Uses

Prosma[®] is indicated for the treatment of allergic conjunctivitis.

Dosage and administration

Adults and children over 3 years of age : The recommended dose is one drop in affected eye(s) 8 to 12 hourly or as recommended by the physician.

Children below 3 years of age: Not recommended.

Contra-indications, warnings, etc.

Prosma[®] is contraindicated in patients with hypersensitivity to any of its ingredient.

Use in Pregnancy and Lactation: There are no clinical trials on the use of **Prosma[®]** in pregnant or nursing women, therefore, **Prosma[®]** should not be used during pregnancy, except if the benefit justifies the potential risk to the fetus.

Warnings: Topical use only. Not for oral use or as injection.

Precautions: As with all ophthalmic preparations containing benzalkonium chloride, patients are advised not to instill **Prosma[®]** (Ketotifen Fumarate ophthalmic solution) while wearing soft (hydrophilic) contact lenses. Wearers of soft contact lenses should be instructed to remove lenses prior to instillation of drops and to wait at least ten minutes after instilling **Prosma[®]** before they insert their contact lenses.

Side effects: Drowsiness and in isolated cases dry mouth and slight dizziness may occur at the beginning of the treatment, but usually disappear spontaneously after a few days.

Directions for use

The following steps should be considered to assure the right use of **Prosma®**.

- a. Proper administration technique for applying the solution to the eye.
- b. Compliance with full course of therapy.
- c. Missed dose:Applying as soon as possible; not applying if almost time for next dose.
- d. Close bottle immediatety after use.
- e. Proper storage of the medicament.

Do not use longer than one month after opening. It is advisable to destroy any remaining medicament after completing treatment.

Pharmaceutical precautions

Store in a cool dry place. Protect from light.

Package quantities

Prosma® Eye Drop: Dropper bottle of 5ml.

® Registered Trade Mark.



ACI Limited
Narayanganj, Bangladesh